

Respiratory sinus arrhythmia and heart period in infancy as correlates of later oppositional defiant and callous-unemotional behaviors

© The Author(s) 2015
Reprints and permissions:
sagepub.co.uk/journalsPermissions.nav
DOI: 10.1177/0165025415605391
ijbd.sagepub.com



Nicholas Wagner,¹ Roger Mills-Koonce,² Michael Willoughby,³
Cathi Propper,⁴ Peter Rehder,² and Noa Gueron-Sela⁴

Abstract

Extant literature suggests that oppositional defiant disorder (ODD) and callous-unemotional (CU) behaviors in childhood and adolescence are associated with distinct patterns of psychophysiological functioning, and that individual differences in these patterns have implications for developmental pathways to disorder. Very little is known about the associations between psychophysiological functioning in infancy and later ODD and CU behaviors. This study examined associations between basal autonomic nervous system (ANS) functioning in infancy and ODD and CU behaviors in later childhood. Using longitudinal heart period (HP) and respiratory sinus arrhythmia (RSA) data from the Durham Child Health and Development Study ($N = 206$), the current study tested associations within a structural equation modeling framework, between continuous measures of HP and RSA across the first 2 years of life, and later ODD and CU behaviors at first grade. Results indicate that ODD and CU behaviors in childhood are associated with lower baseline RSA, but not HP, across infancy. The implications of these findings for developmental models of ODD and CU behaviors are discussed.

Keywords

callous-unemotional behaviors, heart period, oppositional defiant disorder, respiratory sinus arrhythmia

Evidence suggests that callous-unemotional (CU) behaviors, considered to be a downward extension of specific components of adult psychopathy, provide unique insight into disruptive behavior problems in childhood (Frick, Ray, Thornton, & Kahn, 2014a; Wagner, Mills-Koonce, Willoughby, Zvara, & Cox, 2015; Willoughby, Mills-Koonce, Gottfredson, & Wagner, 2014). Often exhibited in the presence of oppositional defiant disorder (ODD) behaviors, CU behaviors describe non-normative emotional, affective, and cognitive deficits—such as a lack of guilt, empathy, and fear, as well as an over-focus on reward and insensitivity to punishment (R. R. Blair, Peschardt, Budhani, Mitchell, & Pine, 2006; Dadds, Fraser, Frost, & Hawes, 2005; Frick & White, 2008; Kotler & McMahon, 2005)—and typically denote greater risk for later antisocial behavior and psychopathy (Lynam, Caspi, Moffitt, Loeber, & Stouthamer-Loeber, 2007; Rowe, Costello, Angold, Copeland, & Maughan, 2010). Research with older children and adolescents suggests that CU behaviors are associated with distinct psychophysiological profiles characterized by reduced baseline functioning and blunted physiological responses to stressors (for a review, see Frick et al., 2014a). Although work investigating the early physiological correlates of later externalizing behavior problems is growing (Beauchaine, 2001; Patriquin, Lorenzi, Scarpa, Calkins, & Bell, 2015), little is known about the early physiological correlates of later CU behaviors. The current study examined associations between baseline respiratory sinus arrhythmia (RSA) and heart period (HP) during infancy and ODD and CU behaviors when children were in first grade.

Early autonomic functioning plays an important role in the emergence of children's self-regulatory capacities (Calkins & Marcovitch, 2010). Theoretical work and extant research suggest that the ability

to maintain homeostasis by shifting attention from internal processes to external demands, a process fundamental to emotional and behavioral adaptation (Calkins, Propper, & Mills-Koonce, 2013), relies heavily on autonomic nervous system (ANS) functioning (Porges, 2007). Heart rate (HR) or heart period (HP; time in between heart beats) are considered broad measures of ANS functioning that reflect both parasympathetic and sympathetic functioning (Dietrich et al., 2007; Stevenson-Hinde & Marshall, 1999; Wagner, Propper, Gueron-Sela, & Mills-Koonce, 2015). There is some evidence that increased baseline HR in middle childhood, marking heightened autonomic functioning, is associated with higher levels of internalizing behavior problems (Dietrich et al., 2007; Monk et al., 2001), and low baseline HR is associated with externalizing problems (Dietrich et al., 2007; Ortiz & Raine, 2004). Although measures of HP may lack specificity given the combined PNS and SNS influences, they provide broad measures of psychophysiological functioning, and therefore have some utility in research that investigates the relationship between autonomic activity and subsequent behavior.

¹ University of North Carolina at Chapel Hill, USA

² University of North Carolina at Greensboro, USA

³ RTI International, Research Triangle Park, NC, USA

⁴ The Center for Developmental Science, Chapel Hill, NC, USA

Corresponding author:

Nicholas Wagner, University of North Carolina at Chapel Hill, CB#3270 Davie Hall, Room 217, Chapel Hill, NC 27599, USA.

Email: wagnern@live.unc.edu

Respiratory sinus arrhythmia (i.e., the variability in heart rate that occurs at the frequency of breathing) is commonly used to index parasympathetic influence on heart-rate variability via the vagus nerve (Porges, 2007). High baseline RSA represents greater myelinated vagal control of the heart, which enables individuals to adapt in the face of a challenge (Calkins et al., 2013; Porges, 1992). Conversely, low baseline RSA indicates reduced myelinated vagal control that may interfere with the ability to regulate behavioral and emotional state (Porges, 2007). Baseline RSA is a correlate of physiological and behavioral adaptation (Porges, Doussard-Roosevelt, Portales, & Suess, 1994; El-Sheikh, 2005; Sulik, Eisenberg, Silva, Spinrad, & Kupfer, 2013) and high levels of baseline RSA are generally associated with positive psychosocial outcomes. For example, a recent prospective study found associations between high baseline RSA in infancy and attenuated stress response, more organized sleep, and better cognitive control at a 10-year follow-up (Feldman, Rosenthal, & Eidelman, 2014). Additionally, high baseline RSA is related to efficient emotion regulation (Beauchaine, 2001; El-Sheikh, 2005; Scarpa, Haden, & Tanaka, 2010), greater cognitive abilities (C. Blair, 2003; Morgan, Aikins, Steffian, Coric, & Southwick, 2007; Watson, Baranek, Roberts, David, & Perryman, 2010), positive social interactions (Patriquin, Lorenzi, Scarpa, & Bell, 2014), and fewer internalizing problems (Neuhaus, Bernier, & Beauchaine, 2014). Low baseline RSA, on the other hand, has been linked to hostility, anxiety disorders, and aggression (for a review, see Beauchaine, 2001).

Limited research on the associations between autonomic functioning and CU behaviors and ODD during later childhood and adolescence suggests that these behaviors are associated with low baseline psychophysiological functioning (Stadler et al., 2008), including low HR (Anastassiou-Hadjicharalambous & Warden, 2008; Dietrich et al., 2007; Frick, Ray, Thornton, & Kahn, 2014b; Raine, 2002), implying that children and adolescents demonstrating high levels of ODD and CU behaviors may exhibit autonomic under-arousal. Provocative etiological models for CU behaviors have drawn from the aforementioned research to posit that low fear and physiological under-arousal early in life may undermine parental socialization efforts (Frick & Morris, 2004) and disrupt processes through which arousal functions to inhibit deviant behavior (Frick et al., 2014b). Specific to parasympathetic functioning, a recent study on adolescents with disruptive behavior problems indicated that the sub-group of children with high CU behaviors had lower basal levels of RSA compared to children without behavior problems (De Wied, van Boxtel, Matthys, & Meeus, 2012), suggesting that these children may have a reduced capacity to organize physiological resources to deal with environmental challenge.

Research on the early psychophysiological correlates of later behavioral problems and CU behaviors is less well established. This is partly due to the fact that few studies have both prospective psychophysiological data from infancy and a sample of children large enough to include adequate numbers with clinically relevant levels of ODD and CU behaviors in later childhood. Important exceptions include recent work by Willoughby, Waschbusch, Moore, and Propper (2011) and Mills-Koonce and colleagues (2015). Using independent samples, each of these studies of infants who later demonstrated clinically relevant levels of ODD and CU behaviors reported findings suggesting associations between CP and CU outcomes and low baseline RSA and HP. Using a clinically-informed grouping approach from the current sample, Willoughby and colleagues (2011) found that infants high on both ODD and CU behaviors at age 36 months demonstrated heightened autonomic

functioning (low HP) during baseline and an aversive social interaction at ages 3 and 6 months. Similarly, in a separate sample Mills-Koonce and colleagues (2015) found that first-grade children showing high levels of conduct problems and CU behaviors had higher levels of autonomic functioning (low baseline HP) and lower parasympathetic functioning (low RSA) at 15 months of age compared to children with low levels of conduct problems and CU behaviors.

Both studies examined the associations between psychophysiological functioning in the first year of life and later ODD and CU behaviors using a clinically informed grouping strategy. Although informed by this work, the current study provides a novel extension of these findings by examining the extent to which RSA and HP functioning, measured across multiple time points in the first 2 years of life, predict continuous measures of later ODD and CU behaviors in a structural equation modeling framework. It is likely that subclinical levels of callous-unemotional behaviors overlap considerably with normative characteristics such as empathic responding, guilt, and conscience (Frick & White, 2008; Knafo, Zahn-Waxler, Van Hulle, Robinson, & Rhee, 2008; Kochanska, 1997). Because of this overlap, and because elevated but subclinical levels of CU behavior pose a risk for child development, much can be gained from studying predictors of variability in continuous measures of these outcomes in a community sample.

The current study

The current study assesses the associations between infants autonomic functioning, measured five times in the first 2 years of life, and continuous measures of ODD and CU behaviors in first grade. Very few studies have examined the associations between psychophysiological functioning in infancy and later ODD and CU behaviors, and no study has done so using multiple measures of psychophysiological functioning across infancy. As such, prior to addressing the primary research aims, the extent to which there is individual variability in growth of HP and RSA across the first 2 years of life was assessed using a latent curve modeling (LCM; Bollen & Curran, 2005) approach, which is subsumed within a broad structural equation modeling framework (SEM; Bollen, 1989) and allows us to examine associations between individual differences in change over time and later ODD and CU outcomes, if such individual variability in growth is evident. Next, the primary goal of this study was to examine the associations between baseline RSA and HP from 3 to 24 months of age and continuous measures of ODD and CU behaviors in first grade using an SEM approach. Given existing findings, we expected that baseline HP and RSA in early life would be negatively associated with ODD and CU behaviors in first grade.

Methods

Participants

The Durham Child Health and Development Study (DCHDS) is a longitudinal study consisting of 206 families whose children were recruited at 3 months of age. The study included only infants who were healthy, full-term, and born without significant complications. Participants were recruited from a primarily urban population living in North Carolina, such that there were equal numbers of African American and European American families, as well as both high- and low-income households. The sample was 57% African American and 43% European American, and approximately 53% of families were low income (below 200% of the

poverty level). The current analyses examine physiological and questionnaire data from home visits that occurred when the infants were aged 3, 6, 12, 18, 24 months, and in first grade. The goals of the current study were addressed using the full sample.

Measures

ODD and CU behaviors. First-grade measures of children's ODD behaviors were derived from the Achenbach System of Empirically Based Assessment (ASEBA; Achenbach & Rescorla, 2004), which was completed by maternal caregivers during a lab visit. The Inventory of Callous Unemotional (ICU; Essau, Sasagawa, & Frick, 2006) traits was used to assess CU behaviors at first grade, providing partial consistency in measurement with the work by Mills-Koonce and colleagues (2015). The ICU was completed by maternal primary caregivers who responded to 24 items on a 4-point Likert scale ranging from 0 (not at all true) to 3 (definitely true). The items that comprise the ICU were developed from other highly established clinical assessments (e.g. APSD, PCL-YV) and include questions about the extent to which the child uses emotions, expresses feelings, cares about getting in trouble, seems cold and uncaring, and hurts others' feelings. The factor structure and predictive utility of the ICU has been confirmed with samples ranging in age from 13 to 20 years of age (see Essau et al., 2006; Fanti, Frick, & Georgiou, 2009; Roose, Bijttebier, Claes, & Lilienfeld, 2011) and with samples as young as age 3 years (see Ezpeleta, de la Osa, Granero, Penelo, & Domènech, 2013).

Heart period and respiratory sinus arrhythmia. As previously mentioned, HP is a measure of the average interbeat interval (IBI; the length of time between heart beats) and is used as an index of broad ANS functioning, whereas RSA is measured as the variation in IBIs linked to respiration and is used as a specific index of parasympathetic functioning of the ANS. The Mini Logger 2000 was used to collect IBIs (Mini Logger 2000; Mini-Mitter Corp., Bend, OR). Researchers placed two electrodes cross-diagonally (i.e., from upper right to lower left) on the child's chest at the beginning of the 3, 6, 12, 18, and 24-month visits for a 2–4-minute measure of baseline cardiac function while at rest. Electrodes were connected to a preamplifier, which transmitted IBIs to a monitor. Porges' (U.S. Patent No. 4,510,944; Porges, 1985) method of calculating RSA and HP, in which a moving polynomial filter is used to remove frequencies lying outside a normal physiological range (0.24 ± 1.04 Hz), was used, and the estimate of RSA is reported in units of $\ln(\text{ms})^2$. The data files were transferred to a computer, and were edited by two reliable researchers using MXEdit software. The two researchers were trained to reliability in MXEdit software (Delta Biometrics, Bethesda, MD) with Porges's Lab at the University of Maryland, and edited the files by visually scanning the data for outlier points relative to adjacent data and replacing those points by dividing them or summing them so that they would be consistent with surrounding data. Consistent with previous work (Willoughby et al., 2011), only participants who had full and sufficient data with less than 10% editing were included in the final dataset. RSA and HP were calculated every 15 seconds for the baseline period using Porges's (1985) method. Larger values of HP indicate lower heart rate and larger values of RSA suggest greater vagal tone.

Additional covariates. *Child's sex* and *race* were collected at the time of recruitment and were determined by the mother or primary caregiver. Infant race and sex was determined by the mother or

primary caregiver. *Family income-to-needs ratio* was determined using the mother's or primary caregiver's report of the total family yearly income at the first grade visit, the size of the family, and the 2003 federal poverty guidelines.

Analytic strategy

Prior to addressing the primary research aims, the extent to which there is individual variability in growth of HP and RSA across the first 2 years of life was assessed using an LCM approach. If such heterogeneity of growth exists, the LCM approach allows for it to be incorporated into subsequent structural models. Next, the primary goals of this study were addressed using a structural equation modeling (SEM; Mplus 5.2, Muthén & Muthén, 2007) approach to examine whether baseline RSA and HP levels in early childhood predict later ODD and CU behaviors in first grade. Model covariates included child's gender, family income-to-needs ratio at first grade, and child's race. Missing data were handled using the full information maximum likelihood methods (Enders & Bandalos, 2001). Overall model fit was determined using root mean square error of approximation (RMSEA), standardized root mean square residual (SRMR), and comparative fit index (CFI). Good fit was defined as CFI values ≥ 0.95 , RMSEA values ≤ 0.06 , and SRMR values ≤ 0.08 (Hu & Bentler, 1999).

Results

Demographics and growth models

Means, standard deviations, and correlations among variables of interest and model covariates are presented in Table 1. Measures of baseline RSA were typically characterized by significant positive correlations ($r = .362-.639, p < .05$) across time points except for the correlation between 12 and 24 months ($r = .095, p = .54$). Measures of baseline HP were characterized by positive correlations across time points that were significant or approached significance ($r = .226-.499, p < .10$). Exceptions include the correlation between 3 and 6 months ($r = .131, p = .25$), the correlation between 3 and 24 months ($r = .137, p = .27$), and the correlation between 6 and 24 months ($r = -.057, p = .72$). Children's sex was not significantly correlated with ODD ($r = -.059, p = .52$) or CU behaviors ($r = .047, p = .60$), but children's race was correlated with income ($r = -.336, p < .01$).

Prior to estimating the full structural model, unconditional latent growth curves were separately estimated for baseline HP and RSA to determine if separate latent variables for HP and for RSA would sufficiently represent early autonomic functioning. Growth curve analyses indicated there was not significant individual variability in the rates of change of baseline RSA or HP over the three time points used in the study ($\eta = 0.021, p = 0.496$ and $\eta = 41.33, p = 0.139$, respectively). Further, there was a non-significant covariance between the intercept and the linear slope components of growth for both RSA and HP, indicating that variability in mean levels of ANS functioning at 3 months were not related to variability in the rate of change. As such, separate latent factors of baseline RSA and HP from 3, 6, 12, 18, and 24 months seem to provide sufficient estimates of the early ANS functioning and are preferred to other analytic approaches (see Bollen, 2002). Kaiser-Meyer-Olkin and Bartlett's test of sphericity supported the factorability of baseline RSA and HP. For the measures of baseline RSA, the Kaiser-Meyer-Olkin measure of sampling adequacy was .75, above the

Table 1. Bivariate correlations between variables of interest and model covariates.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1. Sex (0 = female)	—														
2. Race (0 = White)	-.082	—													
3. Household income	.033	-.336**	—												
4. Baseline respiratory sinus arrhythmia 3m	.078	-.017	.010	—											
5. Baseline respiratory sinus arrhythmia 6m	.019	.134	-.072	.413**	—										
6. Baseline respiratory sinus arrhythmia 12m	-.068	.265**	-.278*	.362**	.639**	—									
7. Baseline respiratory sinus arrhythmia 18m	.001	.283**	-.150	.364**	.425**	.455**	—								
8. Baseline respiratory sinus arrhythmia 24m	.126	.267*	-.113	.213^	.365*	.095	.452**	—							
9. Baseline heart period 3m	.112	-.119	.243*	.609**	.265*	.125	.106	.153	—						
10. Baseline heart period 6m	-.081	.025	.013	.058	.507**	.272^	.019	-.142	.131	—					
11. Baseline heart period 12m	.025	.208*	-.316*	.223*	.322*	.734**	.266*	.017	.315**	.256^	—				
12. Baseline heart period 18m	.148^	.133	-.024	.193^	.338**	.212^	.404**	.311*	.294**	.226^	.392**	—			
13. Baseline heart period 24m	.174	.229*	-.010	.089	.253	.277^	.321*	.382**	.137	-.057	.252^	.499**	—		
14. Callous-unemotional behaviors	.047	-.005	-.147^	-.158	-.134	-.120	-.135	-.114	-.086	.028	-.109	.037	.150	—	
15. Oppositional defiant behaviors	-.059	-.049	-.130	-.310**	-.035	-.211^	-.108	-.055	-.231*	.304*	-.092	.111	-.082	.356**	—
Number	206	206	122	169	96	102	122	79	169	96	102	122	79	122	122
Mean	.51	.57	5.06	3.41	3.66	3.70	4.56	4.82	4.20.1	445.5	450.4	474.2	491.3	.70	1.7
SD	.50	.49	3.54	.98	.88	.99	1.61	1.58	35.02	51.05	36.33	35.36	43.52	.32	2.1
Minimum/Maximum	0/1	0/1	0/21.1	1.1/6.3	1.6/5.7	1.6/6.5	1.6/9.4	1.8/9.6	335.8/520.9	367.5/751.1	373.1/565.5	395.3/568.7	361.4/580.5	.08/1.6	0/9

*p < .05; **p < .01; ^p < .10.

recommended value of .6, and Bartlett's test of sphericity was significant, $X^2(10) = 33.12, p < .01$. For the measures of baseline HP, the Kaiser-Meyer-Olkin measure of sampling adequacy was .62 and Bartlett's test of sphericity was significant, $X^2(10) = 25.43, p < .01$.

Structural equation model

Continuous measures of ODD and CU behavior in first grade were regressed on a latent factor of infant baseline RSA (derived from 3–24 months of age). Lower baseline RSA levels in infancy predicted higher levels of ODD ($\beta = -0.277; p < .05$) and CU ($\beta = -0.251; p < .05$) behaviors in first grade, controlling for race, gender, and family income at first grade. The standardized factor loadings for baseline RSA ranged from .50 to .87 and the final model provided good fit to the data: $X^2(26) = 24.86, p = .52$; CFI = 0.99; RMSEA = 0.001; SRMR = 0.059. Finally, continuous measures of ODD and CU behavior in first grade were regressed on a latent factor of infant baseline HP (derived from 3–24 months of age). These associations were not significant. Standardized coefficients and standard errors for the associations between predictors and outcomes for each model are presented in Table 2.

Discussion

The findings of the current longitudinal study suggest that the early psychophysiological profiles of children who demonstrate high levels of ODD and CU behaviors may be characterized by low baseline levels of RSA in infancy. Associations between baseline HP and ODD and CU behaviors were not observed, suggesting that, in our sample, parasympathetic functioning in the first 2 years of life may be more predictive of later ODD and CU behaviors than a broad measure of ANS functioning. This aligns with findings showing that adolescents high on disruptive and CU behaviors show lower levels of resting RSA, but not HP, than adolescents who demonstrate only disruptive behaviors or neither disruptive nor CU behaviors (De Wied et al., 2012). With regard to infancy, Mills-Koonce and colleagues (2015) observed larger effects for RSA than for HP, which, taken together with the current findings, suggests that specific measures of parasympathetic functioning may be more informative than more broad measures of autonomic functioning for the development of later CU behaviors. These results extend a broader literature on the associations between PNS functioning and the development of disruptive behavioral disorders (Calkins et al., 2013) and ODD and CU behaviors, in particular (De Wied et al., 2012; Mills-Koonce et al., 2015), and advance the literature by demonstrating associations between baseline RSA measured multiple times in the first 2 years of life and continuous measures of later ODD and CU behaviors.

Findings from the current study speak to the importance of considering early baseline physiological functioning in the development of ODD and CU behaviors. Much of the work on the extent to which physiological functioning informs the development of emotion and behavioral regulation has focused on stress physiology (Blair & Raver, 2012) and stress reactivity (Quas et al., 2014). Although the investigation of physiological responses to stress undoubtedly informs etiological models of psychopathology, there is theoretical and empirical support for the utility of examining the associations between baseline physiological functioning and later behavioral problems. For example, it is likely that patterns of stress responses do not become

Table 2. Standardized parameter estimates for the full structural equation model.

	β (B)	95% CI
Respiratory sinus arrhythmia model		
Child gender →	0.07 (0.04)	−0.07 to 0.16
Callous-unemotional behaviors		
Child race →	0.03 (0.02)	−0.11 to 0.16
Callous-unemotional behaviors		
Income first grade →	−0.18* (−0.02)*	−0.03 to 0.00
Callous-unemotional behaviors		
Baseline respiratory sinus arrhythmia →	−0.25* (−0.08)*	−0.14 to −0.01
Callous-unemotional behaviors		
Child gender →	−0.04 (−0.16)	−0.89 to 0.56
Oppositional defiant behaviors		
Child race →	0.00 (−0.01)	−0.87 to 0.87
Oppositional defiant behaviors		
Income first grade →	−0.19* (−0.11)*	−0.22 to −0.01
Oppositional defiant behaviors		
Baseline respiratory sinus arrhythmia →	−0.28* (−0.53)*	−0.98 to −0.08
Oppositional defiant behaviors		
Heart period model		
Child gender →	0.06 (0.04)	−0.08 to 0.16
Callous-unemotional behaviors		
Child race →	−0.05 (−0.03)	−0.16 to 0.09
Callous-unemotional behaviors		
Income first grade →	−0.17 (−0.02)	−0.03 to 0.01
Callous-unemotional behaviors		
Baseline heart period →	−0.04 (−0.01)	−0.09 to 0.06
Callous-unemotional behaviors		
Child gender →	−0.06 (−0.25)	−1.03 to 0.53
Oppositional defiant behaviors		
Child race →	−0.11 (−0.46)	−1.28 to 0.36
Oppositional defiant behaviors		
Income first grade →	−0.16 (−0.10)	−0.20 to 0.01
Oppositional defiant behaviors		
Baseline heart period →	0.01 (0.01)	−0.47 to 0.49
Oppositional defiant behaviors		

* $p < .05$.

integrated until after early developmental periods, suggesting that there is some benefit to restricting initial investigations to baseline measures (Quas et al., 2014). Further, investigations into baseline physiological functioning provide important insight into developmental processes because baseline levels are thought to constrain the magnitude of possible stress response (Cacioppo, Uchino, & Berntson, 1994) and the maintenance and development of baseline physiological functioning in early life is, itself, a dynamic process (Porges, 1992).

There is clear evidence that patterns of RSA inform key regulatory functions implicated in the development and maintenance of pathological behavior patterns (Beauchaine, 2001; Porges, 1996) and the current findings join a body of literature that suggests that the investigation of basal levels of parasympathetic functioning provides important insight into the development of ODD and CU in particular (De Wied et al., 2012; Mills-Koonce et al., 2015; Willoughby et al., 2011). Vagal control of organs including the heart, larynx, and pharynx provides the parasympathetic branch of the ANS substantial control over socially-oriented behaviors such as facial gestures and vocalizations, and consequently contributes to the modulation of social and communicative behaviors (e.g., eye contact, vocal intonation, facial expression). Thus, whereas high vagal control (indicated by high RSA) facilitates sensitive, adaptive

responses to social interactions, low vagal control (low RSA) may be associated with decreased sensitivity and responsiveness to social cues that compromise spontaneous social behavior, social awareness, and affect expressivity (Porges, 2001, 2003, 2007). Consequently, infants' parasympathetic functioning likely plays an important role in establishing mutually positive and responsive interactions with caregivers (Moore et al., 2009; Propper & Moore, 2006).

One possible interpretation of the current findings is that low parasympathetic functioning disrupts early social communication and undermines normative socialization processes (Calkins & Keane, 2004; Calkins & Marcovitch, 2010; Mills-Koonce et al., 2015; Patriquin et al., 2015), setting the stage for the development of ODD and CU behaviors. Indeed, research findings with samples of older children have demonstrated associations between ODD and CU behaviors and behavioral phenotypes that may disrupt the caregiver-child relationship, including deficits in orienting towards caregivers (i.e., less eye contact and mutual orienting) and dysregulated responses to social stimuli (R. R. Blair, 1999; Colder, Mott, & Berman, 2002; Dadds, Jambak, Pasalich, Hawes, & Brennan, 2011; Frick et al., 2003; Frick & White, 2008; Loney, Lima, & Butler, 2006; Mills-Koonce et al., 2014; Willoughby et al., 2011). Early responsive and sensitive interactive behaviors in infancy and early childhood have implications for children's conscience development (e.g., Kochanska, 1997; Kochanska & Murray, 2000) and the development of moral emotion and empathy (Kochanska, Forman, Aksan, & Dunbar, 2005). Further, mutually responsive parent-child interactions have been shown to predict fewer behavior problems for youth high in CU behaviors (Kochanska, Kim, Boldt, & Yoon, 2013). It is likely that future work that examines the associations between early caregiving experiences and autonomic functioning in the prediction of psychopathological outcomes, including ODD and CU behaviors, will be important, particularly considering research indicating that low levels of baseline RSA were associated with greater improvements in child disruptive behavior following a Parent-Child Interaction Therapy (PCIT) program (Bagner et al., 2012).

The strengths of the current study include a prospective longitudinal design, measures of RSA and HP collected at multiple times in early and late infancy, and measures of ODD and CU behaviors collected in first grade. However, these findings should be considered in the context of several limitations. Given the available data, we were only able to investigate differences in baseline RSA and HP functioning. Studies that incorporate measures of both parasympathetic and sympathetic branches of the ANS and examine baseline, reactivity, and recovery of psychophysiological functioning are necessary to provide a more complete picture of how these systems interact to influence development (El-Sheikh, Keiley, Erath, & Dyer, 2013; Nederhof, Marceau, Shirtcliff, Hastings, & Oldehinkel, 2015). Additionally, although the use of a community sample may increase the generalizability of the findings, our ability to examine differences in autonomic functioning between clinically relevant groups was hampered by low sample sizes. With those limitations in mind, the current study replicates a provocative finding regarding the early etiology of a complex and socially significant behavioral phenotype. Future research examining how these early psychophysiological profiles change over time, and how they interact with the caregiving environment (and intervention contexts), are important next steps in identifying the developmental processes underlying these behaviors, as well as informing prevention and treatment.

Acknowledgements

We thank all the parents and children who participated in the Durham Child Health and Development Study and the research assistants for their valuable efforts in collecting this data.

Funding

This study was supported by The North Carolina Child Development Research Collaborative, which is funded by the National Science Foundation through Children's Research Initiative Grant (BCS-0126475). The work was supported in part by a predoctoral fellowship provided by the National Institute of Child Health and Human Development (T32-HD07376) through the Center for Developmental Science, University of North Carolina at Chapel Hill, to Nicholas J. Wagner.

References

- Achenbach, T. M., & Rescorla, L. A. (2004). The Achenbach System of Empirically Based Assessment (ASEBA) for ages 1.5 to 18 years. In M. E. Maruish & M. E. Maruish (Eds.), *The use of psychological testing for treatment planning and outcomes assessment: Volume 2: Instruments for children and adolescents* (3rd ed, pp. 179–213). Mahwah, NJ: Lawrence Erlbaum Associates Publishers.
- Anastassiou-Hadjicharalambous, X., & Warden, D. (2008). Physiologically-indexed and self-perceived affective empathy in Conduct-Disordered children high and low on Callous-Unemotional traits. *Child Psychiatry and Human Development, 39*, 503–517. doi:10.1007/s10578-008-0104-y
- Bagner, D. M., Graziano, P. A., Jaccard, J., Sheinkopf, S. J., Vohr, B. R., & Lester, B. M. (2012). An initial investigation of baseline respiratory sinus arrhythmia as a moderator of treatment outcome for young children born premature with externalizing behavior problems. *Behavior Therapy, 43*, 652–665. doi:10.1016/j.beth.2011.12.002
- Beauchaine, T. (2001). Vagal tone, development, and Gray's motivational theory: Toward an integrated model of autonomic nervous system functioning in psychopathology. *Development and Psychopathology, 13*, 183–214. doi:10.1017/S0954579401002012
- Blair, C. (2003). Behavioral inhibition and behavioral activation in young children: Relations with self-regulation and adaptation to preschool in children attending head start. *Developmental Psychobiology, 42*, 301–311. doi:10.1002/dev.10103
- Blair, C., & Raver, C. C. (2012). Individual development and evolution: Experiential canalization of self-regulation. *Developmental Psychology, 48*, 647–657. doi:10.1037/a0026472
- Blair, R. R. (1999). Responsiveness to distress cues in the child with psychopathic tendencies. *Personality and Individual Differences, 27*, 135–145. doi:10.1016/S0191-8869(98)00231-1
- Blair, R. R., Peschardt, K., Budhani, S., Mitchell, D. V., & Pine, D. S. (2006). The development of psychopathy. *Journal of Child Psychology and Psychiatry, 47*, 262–275. doi:10.1111/j.1469-7610.2006.01596.x
- Bollen, K. A. (1989). *Structural equations with latent variables*. Series in probability and mathematical statistics. Hoboken, NJ: Wiley & Sons.
- Bollen, K. A. (2002). Latent variables in psychology and the social sciences. *Annual Review of Psychology, 53*(1), 605–634. doi:10.1146/annurev.psych.53.100901.135239
- Bollen, K. A., & Curran, P. J. (2006). *Latent curve models: A structural equation perspective*. Hoboken, NJ: Wiley & Sons.

- Cacioppo, J. T., Uchino, B. N., & Berntson, G. G. (1994). Individual differences in the autonomic origins of heart rate reactivity: The psychometrics of respiratory sinus arrhythmia and preejection period. *Psychophysiology*, *31*, 412–419. doi:10.1111/j.1469-8986.1994.tb02449.x
- Calkins, S. D., & Keane, S. P. (2004). Cardiac vagal regulation across the preschool period: Stability, continuity, and implications for childhood adjustment. *Developmental Psychobiology*, *45*, 101–112. doi:10.1002/dev.20020
- Calkins, S. D., & Marcovitch, S. (2010). Emotion regulation and executive functioning in early development: Integrated mechanisms of control supporting adaptive functioning. In S. D. Calkins, M. A. Bell, S. D. Calkins & M. A. Bell (Eds.), *Child development at the intersection of emotion and cognition* (pp. 37–57). Washington, DC: American Psychological Association. doi:10.1037/12059-003
- Calkins, S. D., Propper, C., & Mills-Koonce, W. R. (2013). A biosychosocial perspective on parenting and developmental psychopathology. *Development and Psychopathology*, *25*, 1399–1414. doi:10.1017/S0954579413000680
- Colder, C. R., Mott, J. A., & Berman, A. S. (2002). The interactive effects of infant activity level and fear on growth trajectories of early childhood behavior problems. *Development and Psychopathology*, *14*, 1–23. doi:10.1017/S0954579402001013
- Dadds, M. R., Fraser, J., Frost, A., & Hawes, D. J. (2005). Disentangling the underlying dimensions of psychopathy and conduct problems in childhood: A community study. *Journal of Consulting and Clinical Psychology*, *73*, 400–410. doi:10.1037/0022-006X.73.3.400
- Dadds, M. R., Jambak, J., Pasalich, D., Hawes, D. J., & Brennan, J. (2011). Impaired attention to the eyes of attachment figures and the developmental origins of psychopathy. *Journal of Child Psychology and Psychiatry*, *52*, 238–245. doi:10.1111/j.1469-7610.2010.02323.x
- De Wied, M., Van Boxtel, A., Matthys, W., & Meeus, W. (2012). Verbal, facial and autonomic responses to empathy-eliciting film clips by disruptive male adolescents with high versus low callous-unemotional traits. *Journal of Abnormal Child Psychology*, *40*, 211–223. doi:10.1007/s10802-011-9557-8
- Dietrich, A., Riese, H., Sondejker, F. E. P. L., Greaves-Lord, K., van Roon, A. M., Ormel, J., . . . Rosmalen, J. G. M. (2007). Externalizing and internalizing problems in relation to autonomic function: A population-based study in preadolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, *46*, 378–386. doi:10.1097/CHI.0b013e31802b91ea
- El-Sheikh, M. (2005). Stability of respiratory sinus arrhythmia in children and young adolescents: A longitudinal examination. *Developmental Psychobiology*, *46*, 66–74. doi:10.1002/dev.20036
- El-Sheikh, M., Keiley, M., Erath, S., & Dyer, W. J. (2013). Marital conflict and growth in children's internalizing symptoms: The role of autonomic nervous system activity. *Developmental Psychology*, *49*, 92–108. doi:10.1037/a0027703
- Enders, C. K., & Bandalos, D. L. (2001). The relative performance of full information maximum likelihood estimation for missing data in structural equation models. *Structural Equation Modeling*, *8*, 430–457. doi:10.1207/S15328007sem0803_5
- Essau, C. A., Sasagawa, S., & Frick, P. J. (2006). Callous-unemotional traits in a community sample of adolescents. *Assessment*, *13*, 454–469. doi:10.1177/1073191106287354
- Ezpeleta, L., de la Osa, N., Granero, R., Penelo, E., & Domènech, J. M. (2013). Inventory of callous-unemotional traits in a community sample of preschoolers. *Journal of Clinical Child and Adolescent Psychology*, *42*, 91–105. doi:10.1080/15374416.2012.734221
- Fanti, K. A., Frick, P. J., & Georgiou, S. (2009). Linking callous-unemotional traits to instrumental and non-instrumental forms of aggression. *Journal of Psychopathology and Behavioral Assessment*, *31*, 285–298. doi:10.1007/s10862-008-9111-3
- Feldman, R., Rosenthal, Z., & Eidelman, A. I. (2014). Maternal-preterm skin-to-skin contact enhances child physiologic organization and cognitive control across the first 10 years of life. *Biological Psychiatry*, *75*, 56–64. doi:10.1016/j.biopsych.2013.08.012
- Frick, P. J., Cornell, A. H., Bodin, S. D., Dane, H. E., Barry, C. T., & Loney, B. R. (2003). Callous-unemotional traits and developmental pathways to severe conduct problems. *Developmental Psychology*, *39*(2), 246–260. doi:10.1037/0012-1649.39.2.246
- Frick, P. J., & Morris, A. S. (2004). Temperament and developmental pathways to conduct problems. *Journal of Clinical Child and Adolescent Psychology*, *33*, 54–68. doi:10.1207/S15374424JCCP3301_6
- Frick, P. J., Ray, J. V., Thornton, L. C., & Kahn, R. E. (2014a). Annual research review: A developmental psychopathology approach to understanding callous-unemotional traits in children and adolescents with serious conduct problems. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, *55*, 532–48. doi:10.1111/jcpp.12152
- Frick, P. J., Ray, J. V., Thornton, L. C., & Kahn, R. E. (2014b). Can callous-unemotional traits enhance the understanding, diagnosis, and treatment of serious conduct problems in children and adolescents? A comprehensive review. *Psychological Bulletin*, *140*, 1–57. doi:10.1037/a0033076
- Frick, P. J., & White, S. F. (2008). Research review: The importance of callous-unemotional traits for developmental models of aggressive and antisocial behavior. *Journal of Child Psychology and Psychiatry*, *49*, 359–375. doi:10.1111/j.1469-7610.2007.01862.x
- Hu, L. T., & Bentler, P. M. (1999). Cutoff criteria for fit indices in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling*, *6*, 1–55. doi:10.1080/10705519909540118
- Knafo, A., Zahn-Waxler, C., Van Hulle, C., Robinson, J. L., & Rhee, S. H. (2008). The developmental origins of a disposition toward empathy: Genetic and environmental contributions. *Emotion*, *8*, 737–752. doi:10.1037/a0014179
- Kochanska, G. (1997). Multiple pathways to conscience for children with different temperaments: From toddlerhood to age 5. *Developmental Psychology*, *33*, 228–240. doi:10.1037/0012-1649.33.2.228
- Kochanska, G., Forman, D. R., Aksan, N., & Dunbar, S. B. (2005). Pathways to conscience: Early mother-child mutually responsive orientation and children's moral emotion, conduct, and cognition. *Journal of Child Psychology and Psychiatry*, *46*, 19–34. doi:10.1111/j.1469-7610.2004.00348.x
- Kochanska, G., Kim, S., Boldt, L. J., & Yoon, J. E. (2013). Children's callous-unemotional traits moderate links between their positive relationships with parents at preschool age and externalizing behavior problems at early school age. *Journal of Child Psychology and Psychiatry*, *54*, 1251–1260.
- Kochanska, G., & Murray, K. T. (2000). Mother-child mutually responsive orientation and conscience development: From toddler

- to early school age. *Child Development*, 71, 417–431. doi:10.1111/1467-8624.00154
- Kotler, J. S., & McMahon, R. J. (2005). Child psychopathy: Theories, measurement, and relations with the development and persistence of conduct problems. *Clinical Child and Family Psychology Review*, 8, 291–325. doi:10.1007/s10567-005-8810-5
- Loney, B. R., Lima, E. N., & Butler, M. A. (2006). Trait affectivity and nonreferred adolescent conduct problems. *Journal of Clinical Child and Adolescent Psychology*, 35, 329–336. doi:10.1207/s15374424jccp3502_17
- Lynam, D. R., Caspi, A., Moffitt, T. E., Loeber, R., & Stouthamer-Loeber, M. (2007). Longitudinal evidence that psychopathy scores in early adolescence predict adult psychopathy. *Journal of Abnormal Psychology*, 116, 155–165. doi:10.1037/0021-843X.116.1.155
- Mills-Koonce, W. R., Wagner, N. J., Willoughby, M. T., Stifter, C., Blair, C., & Granger, D. A. (2015). Greater fear reactivity and psychophysiological hyperactivity among infants with later conduct problems and callous-unemotional traits. *Journal of Child Psychology and Psychiatry*, 56, 147–154. doi:10.1111/jcpp.12289
- Monk, C., Kovelenco, P., Ellman, L. M., Sloan, R. P., Bagiella, E., Gorman, J. M., & Pine, D. S. (2001). Enhanced stress reactivity in paediatric anxiety disorders: Implications for future cardiovascular health. *International Journal of Neuropsychopharmacology*, 4, 199–206. doi:10.1017/S146114570100236X
- Moore, G. A., Hill-Soderlund, A. L., Propper, C. B., Calkins, S. D., Mills-Koonce, W. R., & Cox, M. J. (2009). Mother–infant vagal regulation in the face-to-face still-face paradigm is moderated by maternal sensitivity. *Child Development*, 80(1), 209–223. doi:10.1111/j.1467-8624.2008.01255.x
- Morgan, C. A., Aikins, D. E., Steffian, G., Coric, V., & Southwick, S. (2007). Relation between cardiac vagal tone and performance in male military personnel exposed to high stress: Three prospective studies. *Psychophysiology*, 44, 120–127. doi:10.1111/j.1469-8986.2006.00475.x
- Muthén, L. K., & Muthén, B. O. (2007). *Mplus User's Guide* (6th Ed.). Los Angeles, CA: Muthén & Muthén.
- Nederhof, E., Marceau, K., Shirtcliff, E. A., Hastings, P. D., & Oldehinkel, A. J. (2015). Autonomic and adrenocortical interactions predict mental health in late adolescence: The TRAILS Study. *Journal of Abnormal Child Psychology*, 43(5), 847–861. doi:10.1007/s10802-014-9958-6
- Neuhaus, E., Bernier, R., & Beauchaine, T. P. (2014). Brief report: Social skills, internalizing and externalizing symptoms, and respiratory sinus arrhythmia in autism. *Journal of Autism and Developmental Disorders*, 44, 730–737. doi:10.1007/s10803-013-1923-7
- Ortiz, J., & Raine, A. (2004). Heart rate level and antisocial behavior in children and adolescents: A meta-analysis. *Journal of the American Academy of Child & Adolescent Psychiatry*, 43, 154–162. doi:10.1097/00004583-200402000-00010
- Patriquin, M. A., Lorenzi, J., Scarpa, A., & Bell, M. A. (2014). Developmental trajectories of respiratory sinus arrhythmia: Associations with social responsiveness. *Developmental Psychobiology*, 56, 317–326. doi:10.1002/dev.21100
- Patriquin, M. A., Lorenzi, J., Scarpa, A., Calkins, S. D., & Bell, M. A. (2015). Broad implications for respiratory sinus arrhythmia development: Associations with childhood symptoms of psychopathology in a community sample. *Developmental Psychobiology*, 57, 120–130. doi:10.1002/dev.21269
- Porges, S. W. (1985). Method and apparatus for evaluating rhythmic oscillations in a periodic physiological response system (U.S. Patent No. 4,510,944). Washington, DC: U.S. Patent and Trademark Office.
- Porges, S. W. (1992). Vagal tone: A physiologic marker of stress vulnerability. *Pediatrics*, 90, 498–504.
- Porges, S. W. (1996). Physiological regulation in high-risk infants: A model for assessment and potential intervention. *Development and Psychopathology*, 8, 29–42. doi:10.1017/S0954579400006969
- Porges, S. W. (2001). The polyvagal theory: Phylogenetic substrates of a social nervous system. *International Journal of Psychophysiology*, 42, 123–146. doi:10.1016/S0167-8760(01)00162-3
- Porges, S. W. (2003). The polyvagal theory: Phylogenetic contributions to social behavior. *Physiology & Behavior*, 79, 503–513. doi:10.1016/S0031-9384(03)00156-2
- Porges, S. W. (2007). The polyvagal perspective. *Biological Psychology*, 74, 116–143. doi:10.1016/j.biopsycho.2006.06.009
- Porges, S. W., Doussard-Roosevelt, J. A., Portales, L. A., & Suess, P. E. (1994). Cardiac vagal tone: Stability and relation to difficulty in infants and 3-year-olds. *Developmental Psychobiology*, 27, 289–300. doi:10.1002/dev.420270504
- Propper, C., & Moore, G. A. (2006). The influence of parenting on infant emotionality: A multi-level psychobiological perspective. *Developmental Review*, 26, 427–460. doi:10.1016/j.dr.2006.06.003
- Quas, J. A., Yim, I. S., Oberlander, T. F., Nordstokke, D., Essex, M. J., Armstrong, J. M., . . . Boyce, W. T. (2014). The symphonic structure of childhood stress reactivity: Patterns of sympathetic, parasympathetic, and adrenocortical responses to psychological challenge. *Development and Psychopathology*, 26, 963–982. doi:10.1017/S0954579414000480
- Raine, A. (2002). Biosocial studies of antisocial and violent behavior in children and adults: A review. *Journal of Abnormal Child Psychology*, 30, 311–326. doi:10.1023/A:1015754122318
- Roose, A., Bijttebier, P., Claes, L., & Lilenfeld, S. O. (2011). Psychopathic traits in adolescence: Associations with the revised Reinforcement Sensitivity Theory systems. *Personality and Individual Differences*, 50, 201–205. doi:10.1016/j.paid.2010.09.028
- Rowe, R., Costello, E. J., Angold, A., Copeland, W. E., & Maughan, B. (2010). Developmental pathways in oppositional defiant disorder and conduct disorder. *Journal of Abnormal Psychology*, 119, 726–738. doi:10.1037/a0020798
- Scarpa, A., Haden, S. C., & Tanaka, A. (2010). Being hot-tempered: Autonomic, emotional, and behavioral distinctions between childhood reactive and proactive aggression. *Biological Psychology*, 84, 488–496. doi:10.1016/j.biopsycho.2009.11.006
- Stadler, C., Grasmann, D., Fegert, J. M., Holtmann, M., Poustka, F., & Schmeck, K. (2008). Heart rate and treatment effect in children with disruptive behavior disorders. *Child Psychiatry and Human Development*, 39(3), 299–309. doi:10.1007/s10578-007-0089-y
- Stevenson-Hinde, J., & Marshall, P. J. (1999). Behavioral inhibition, heart period, and respiratory sinus arrhythmia: An attachment perspective. *Child Development*, 70, 805–816. doi:10.1111/1467-8624.00058
- Sulik, M. J., Eisenberg, N., Silva, K. M., Spinrad, T. L., & Kupfer, A. (2013). Respiratory sinus arrhythmia, shyness, and effortful control in preschool-age children. *Biological Psychology*, 92, 241–248. doi:10.1016/j.biopsycho.2012.10.009

- Wagner, N. J., Mills-Koonce, W. R., Willoughby, M. T., Zvara, B., & Cox, M. J. (2015). Parenting and children's representations of family predict disruptive and callous-unemotional behaviors. *Developmental Psychology, 51*, 935–948. doi:10.1037/a0039353
- Wagner, N. J., Propper, C., Gueron-Sela, N., & Mills-Koonce, W. R. (2015). Dimensions of maternal parenting and infants' autonomic functioning interactively predict early internalizing behavior problems. *Journal of Abnormal Child Psychology*. Advance online publication. doi:10.1007/s10802-015-0039-2
- Watson, L. R., Baranek, G. T., Roberts, J. E., David, F. J., & Perryman, T. Y. (2010). Behavioral and physiological responses to child-directed speech as predictors of communication outcomes in children with autism spectrum disorders. *Journal of Speech, Language, and Hearing Research, 53*, 1052–1064. doi:10.1044/1092-4388(2009/09-0096)
- Willoughby, M. T., Mills-Koonce, W. R., Gottfredson, N. C., & Wagner, N. J. (2014). Measuring callous unemotional behaviors in early childhood: Factor structure and the prediction of stable aggression in middle childhood. *Journal of Psychopathology and Behavioral Assessment, 36*, 30–42. doi:10.1007/s10862-013-9379-9
- Willoughby, M. T., Waschbusch, D. A., Moore, G. A., & Propper, C. B. (2011). Using the ASEBA to screen for callous unemotional traits in early childhood: Factor structure, temporal stability, and utility. *Journal of Psychopathology and Behavioral Assessment, 33*, 19–30. doi:10.1007/s10862-010-9195-4